

CHECKMATE[®] OFM Dispenser

Active Ingredients:

(Z)-8-Dodecen-1-yl acetate 11.93%
(E)-8-Dodecen-1-yl acetate.....0.80%
(Z)-8-Dodecen-1-ol0.15%
Other Ingredients: 87.12%
Total:100.00%

Net Weight:

250 mg active ingredients per dispenser.



**A BIOCHEMICAL PESTICIDE FOR MATING DISRUPTION OF
ORIENTAL FRUIT MOTH (*Grapholita molesta*), MACADAMIA NUT BORER
(*Cryptophlebia ombrodelta*), and KOA SEEDWORM (*Cryptophlebia illepipa*)**

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid contact of pheromone reservoir inside dispenser with skin or clothing. Wash thoroughly with soap and water after handling.

FIRST AID STATEMENT

IF ON SKIN: In case of contact, flush area with water and wash thoroughly with soap and water. Obtain medical attention if irritation persists.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

HOT LINE NUMBER: Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-866-326-6737 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by disposal of wastes.

WARRANTY AND LIMITATION OF DAMAGES

Suterra, LLC warrants that this material conforms to the chemical description on the label. **Suterra** neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, expressed or implied, concerning this material. **Suterra's** maximum liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this material.

Lot #:

Operator: **Shift: D S**

Machine: **Bucket:**

NET CONTENTS:

DISPENSERS

OFM

Dispenser

8

1

0

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CheckMate® OFM Dispenser is a behavior modifying biochemical product (pheromone) which functions by mating disruption. It is to be used in pest management programs for control of oriental fruit moth (*Grapholita molesta*), macadamia nut borer (*Cryptophlebia ombrodelta*), and koa seedworm (*Cryptophlebia illepidata*).

METHOD OF APPLICATION: For use on peaches, nectarines, apricots, plums, cherries and other stone fruits; apples, quince and other pome fruits; almonds, macadamia and other tree nut crops. The rate of application of **CheckMate® OFM Dispenser** is 100-200 dispensers per acre. Apply dispensers by hand to thumb-size limbs or hangers of crop to be protected. Place dispensers in the upper one-third of the tree, or higher. If there is a persistent and prevalent wind, apply one dispenser to every tree on the upwind side of the orchard and then evenly distribute the remaining dispensers in the block. **Do not exceed a total of 150 grams a.i./acre/year.**

Wash hands after application. Do not handle other types of insect lures before or after lure placement without thoroughly washing hands. Place monitoring traps in the orchard before handling **CheckMate® OFM Dispenser**. Monitor treatments carefully, especially for secondary pests such as peach twig borer.

Reseal containers immediately after use. Store unused product in a refrigerator or freezer not used for food storage. Do not use **CheckMate® OFM Dispenser** from damaged, punctured or unsealed containers.

RECOMMENDED APPLICATION: For best results, apply at the time or shortly before oriental fruit moths, macadamia nut borer moths or koa seedworm moths begin to emerge in the spring (biofix), or when the moths are first detected in pheromone traps such as **BioLure®**. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the orchard must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection to developing fruit. These supplemental insecticide applications should be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control emerging larvae. Follow the **BioLure® Technical Bulletin** for trap placement. Monitor for insect infestation with traps and by visual inspection of trees and fruit. Re-apply **CheckMate® OFM Dispenser** as needed, based on monitoring results and field scouting.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE - Store container in cool place until used.

PESTICIDE DISPOSAL - Wastes resulting from the use of this product may be disposed of on-site or at an approved disposal facility.

CONTAINER DISPOSAL - Do not reuse container. Discard in trash.

Manufactured by:

SUTERRA™

Suterra, LLC
213 Southwest Columbia St.
Bend, Oregon, USA 97702
1-866-326-6737
Fax (541) 388-3705

EPA Registration No. 56336-36
EPA Establishment No. 56336-OR-1
ofmdiseng810-04 12/2002